



POLICY BRIEF

CUHAS/PB/002/2017

Introduction of congenital viral infections surveillance and screening services for pregnant women in Tanzania

Key messages

- Viral infections during pregnancy are common in Tanzania and have been found to cause inborn (congenital) abnormalities
- About 36.5% of pregnant women in Tanzania are infected with at least one virus that is capable of causing inborn abnormalities
- Despite all these information, there are no specific preventive strategies to address congenital virus infections in Tanzania.
- This policy brief recommends the introduction of surveillance and screening services for viral infections among pregnant women in Tanzania

Executive summary

Despite high percentage of children born with inborn abnormalities, Tanzania has no birth defects registry to provide information on the extent, trends and patterns of the problem. A large proportion of pregnant women in Tanzania is infected with viruses such as Rubella, Cytomegalovirus (CMV), Parvovirus B19 (B19) and Hepatitis B that are known to cause inborn abnormalities. In these cases, undesirable pregnancy outcomes such as spontaneous abortions, stillbirths and congenital abnormalities have been reported. The impact associated with these infections is lifetime health costs related to treatment and caring of the affected individuals as well as socioeconomic and social stress to the family and community. Despite all these information, there are no specific preventive strategies to address congenital virus infections in Tanzania. This policy brief recommends the

introduction of surveillance and screening services for inborn viral infections among pregnant women in Tanzania. Such services will help to establish risk transmission levels of these viruses to the unborn babies and guide on preventive measures including alerts for possible future outbreaks in the country.

The Problem

Despite the fact that inborn abnormalities are common in Tanzania (Mashuda et al., 2014) and might be due to various infections, the current practice regarding screening during pregnancy only targets Syphilis and Human immunodeficiency virus (HIV). Other infections such as of *Toxoplasma gondii*, Parvovirus B19 (B19), Hepatitis B virus (HBV), Rubella virus, Cytomegalovirus (CMV), Herpes simplex type 2 virus (HSV-2), Chikungunya virus and Zika virus that are also known to cause inborn abnormalities/infections, that are endemic in the country, are not usually screened at health facilities. Unfortunately, Tanzania does not have birth defect registry which could be used to track the trends and patterns of inborn abnormalities for possible and appropriate interventions.

Studies in Tanzania have shown that about 36.5% of pregnant women are infected with at least one virus that is capable of causing inborn infections and more than one third of these women have indicators showing that they have been previously infected with more than one inborn viral infection (Rubella, CMV and Parvovirus B19) (Mhalu & Haukenes, 1990; Mwambe et al., 2014; Mirambo et al., 2016a). In addition, studies from Dar es Salaam and Mwanza have shown that percentage of pregnant women infected with Hepatitis B virus ranges from 2 to 7% (Rashid et al., 2014; Mirambo et al., 2016b) which is considered as relative

higher (WHO, 2009). More studies in Tanzania have shown that about 1,000 out of 100,000 of the live new-borns are likely to have congenital rubella syndrome (CRS) and about 4 in every 1,000 women are likely to deliver babies with inborn CMV infections. These statistics indicate that viral infections are common and are strongly associated with poor pregnancy outcomes (Chibwe, 2016; Mirambo et al., 2016c). In addition Tanzania has been classified as endemic country to Zika virus infections (<https://wwwnc.cdc.gov/>), an infection first reported in the country in 1952. Some cases of Chikungunya virus infections have been reported in Tanzania (Hertz et al., 2012; Chipwaza et al., 2014; Kajeguka et al., 2016). Zika and Chikungunya viruses have been confirmed to cause brain and other inborn abnormalities in studies elsewhere (Gérardin et al., 2014; Sarno et al., 2016; Valentine et al., 2016; van der Linden et al., 2016).

Inborn abnormalities are associated with lifetime health costs, socioeconomic impact and social stress to the family and community. Based on the records on CRS, the lifetime treatment can cost US\$14,000-200,000 per individual patient (Irons et al., 2000; Duszak, 2009; Castillo-Solórzano et al., 2003) which is very high for a country like Tanzania. This policy brief highlights the importance of considering screening of viral infections which appear to be common across the country.

Policy options

Despite the fact that HIV infection and syphilis are screened during pregnancy, and there is immunization for hepatitis B and Rubella viruses among children under five years of age, it is important to introduce screening for other inborn viral infections for evidence based counselling and appropriate management. Information generated through screening will strengthen our knowledge on the magnitude and trends of these infections for the possible future outbreaks. The following policy options are recommended:

1. **Introduction of screening inborn viruses during pregnancy:** There is evidence that Rubella, CMV, Chikungunya, Hepatitis B and Herpes simplex virus infections are common in Tanzania and might contribute extensively to the cases of inborn abnormalities. Screening for these viruses will be one step ahead in

controlling impacts related to these infections across the country.

2. **Establish a birth defect registry and screening every child with inborn abnormalities:** In order to have a true picture of the impact of these infections, it is important to establish a birth defect registry. It is also important to screen every child with inborn abnormality for the purpose of monitoring of inborn acquired infections.
3. **Introduce evidence based counselling to prevent infections and undesirable pregnancy outcomes:** Having a child with inborn abnormalities is associated with socioeconomic and social stress impacts. To lessen fear among pregnant women and families especially with the current Zika problem worldwide, there is a need to introduce evidence based counselling regarding pregnancy outcome in relation to possible inborn infections.

Implementation considerations

Point of care rapid tests are available for detection of Herpes simplex type 2 virus, Hepatitis B virus, Rubella virus, Cytomegalovirus, Parvovirus B19 justifying the easy implementations of the proposed policy options. These tests do not require advanced skills and infrastructures that would probably require capital investment and maintenance costs. In addition, there are other reliable advanced tests (immunoassays) for Zika virus detection (Huzly et al., 2016). The cost of screening one viral infection using disposable rapid test is about US\$ 1 - there are expected additional costs using this strategy. The current screening services during pregnancy involve drawing of blood for blood typing, level of haemoglobin, syphilis and HIV testing. The same blood sample can be used to screen for other inborn viral infections. Using existing services during pregnancy the identified viral infections can be screened for every pregnant woman attending antenatal clinic. In cases where the test is not available the samples can be taken to the nearby zonal hospitals/health centres for processing using the existing infrastructures. It is important to note that the implementation of these policy options will save the Government and communities from incurring high lifetime

treatment costs (>US\$ 14,000) and social stress associated with the conditions.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

The authors acknowledge support from members of the Department of Microbiology and Immunology-CUHAS, Department of Paediatric

Child health-CUHAS, Department of Obstetrics and Gynaecology-CUHAS, Department of Microbiology and Immunology-MUHAS and the National institute for Medical research. The development of this policy brief was financially supported by the Skoll Global Threats Fund through Enhancing Community-based Disease Outbreak Detection and Response project, Kilimanjaro Christian Medical University College and the National Health Policy and Systems Research Hub.

References

- Castillo-Solórzano, C., Carrasco, P., Tambini, G., Reef, S. et al. (2003) New horizons in the control of rubella and prevention of congenital rubella syndrome in the Americas. *Journal of Infectious Diseases* 187(Suppl. 1), S146-S152.
- Chibwe, E. (2016) Seroprevalence and associated factors of human cytomegalovirus among pregnant women attending antenatal clinics in Mwanza city, Tanzania. MMed Dissertation. *Catholic University of Health and Allied Sciences (CUHAS-Bugando)*.
- Chipwaza, B., Mugasa, J.P., Selemani, M., Amuri, M. et al. (2014) Dengue and Chikungunya fever among viral diseases in outpatient febrile children in Kilosa District Hospital, Tanzania. *PLoS Neglected Tropical Diseases* 8(11): e3335.
- Duszak, R. S. (2009) Congenital rubella syndrome-major review. *Optometry-Journal of the American Optometric Association* 80: 36-43.
- Gérardin, P., Sampéris, S., Ramful, D., Boumahni, B. et al. (2014) Neurocognitive outcome of children exposed to perinatal mother-to-child Chikungunya virus infection: the CHIMERE cohort study on Reunion Island. *PLoS Neglected Tropical Disease* 8(7): e2996.
- Hertz, J.T., Munishi, O.M., Ooi, E.E., Howe, S. et al. (2012) Chikungunya and Dengue Fever among Hospitalized Febrile Patients in Northern Tanzania. *American Journal of Tropical Medicine Hygiene* 86: 171-177.
- Huzly, D., Hanselmann, I., Schmidt-Chanasit, J. & Panning, M. (2016) High specificity of a novel Zika virus ELISA in European patients after exposure to different flaviviruses. *European Surveillance* 21: 30203.
- Irons, B., Lewis, M.J., Dahl-Regis, M., Castillo-Solórzano, C. et al. (2000) Strategies to eradicate rubella in the English-speaking Caribbean. *American Journal of Public Health* 90: 1545.
- Kajeguka, D.C., Kaaya, R.D., Mwakalinga, S., Ndossi, R., et al., (2016) Prevalence of dengue and chikungunya virus infections in north-eastern Tanzania: a cross sectional study among participants presenting with malaria-like symptoms. *BMC Infectious Diseases* 16:183.
- Mashuda, F., Zuechner, A., Challya, P., Kidenya, B.R. et al. (2014) Pattern and factors associated with congenital anomalies among young infants admitted at Bugando medical centre, Mwanza, Tanzania. *BMC Research Notes* 7(1): 195.
- Mhalu, F. & Haukenes, G. (1990) Prevalence of cytomegalovirus antibody in pregnant women, AIDS patients and STD patients in Dar es Salaam. *AIDS* 4, 1294.
- Mirambo, M.M., Aboud, S., Mushi, M.F., Seugendo, M. et al. (2016a) Serological evidence of acute rubella infection among under-fives in Mwanza: a threat to increasing rates of congenital rubella syndrome in Tanzania. *Italian Journal of Pediatrics* 42(1), 1.
- Mirambo, M.M., Chibwe, E., Mushi, M.F., Majigo, M. et al. (2016b) Cytomegalovirus, Parvovirus B19 and Rubella co-infection among pregnant women attending antenatal clinics in Mwanza City: the need to be considered in Tanzanian antenatal care package. *Epidemiology: Open Access*, 2016.

- Mirambo, M.M., Matemba, L., Majigo, M. & Mshana, S.E. (2016c) Zika virus infections among neonates with congenital anomalies. Weill Bugando School of Medicine. *Unpublished Research project*.
- Mirambo, M.M., Mbena, P.B., Mushi, M.F., Mtebe, M. et al. (2016d) Prevalence of Hepatitis B surface antigen among pregnant women attending antenatal clinic at Nyamagana District Hospital Mwanza, Tanzania. *Tanzania Journal of Health Research* 18(1).
- Mwambe, B., Mirambo, M. M., Mshana, S. E., Massinde, A.N. et al. (2014) Sero-positivity rate of rubella and associated factors among pregnant women attending antenatal care in Mwanza, Tanzania. *BMC Pregnancy and Childbirth* 14: 95.
- Rashid, S., Kilewo, C. & Aboud, S. (2014) Seroprevalence of hepatitis B virus infection among antenatal clinic attendees at a tertiary hospital in Dar es Salaam, Tanzania. *Tanzania Journal of Health Research* 16 (1).
- Sarno, M., Sacramento, G. A., Khouri, R., do Rosário, M.S., et al. (2016) Zika virus infection and stillbirths: a case of hydrops fetalis, hydranencephaly and fetal demise. *PLoS Neglected Tropical Disease* 10: e0004517.
- Valentine, G., Marquez, L. & Pammi, M. (2016) Zika virus-associated microcephaly and eye lesions in the newborn. *Journal of the Pediatric Infectious Diseases Society* 5: 323-328.
- van der Linden, V., Rolim Filho, E. L., Lins, O. G., van der Linden, A. et al. (2016) Congenital Zika syndrome with arthrogryposis: retrospective case series study. *BMJ* 354: i3899.
- WHO (2009) World Health Organization Hepatitis B position statement. *Weekly Epidemiological Records* 84: 405–420

About the authors

Mariam M. Mirambo¹, Elieza Chibwe², Lucas Matemba³, Majigo Mtebe⁴, Leonard E.G. Mboera³ & Stephen E. Mshana¹

¹ Department of Microbiology and Immunology, Weill Bugando School of Medicine, Catholic University of Health and Allied sciences, P.O Box 1464, Mwanza, Tanzania

² Department of Obstetrics and Gynaecology, Weill Bugando School of Medicine, Catholic University of Health and Allied Sciences, P.O Box 1464, Mwanza, Tanzania

³ National Institute for Medical Research, P.O. Box 9653, Dar es Salaam, Tanzania

⁴ Department of Microbiology and Immunology, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania

Correspondence: Eng. Eric Beda Mutagahywa; E-mail: eric.beda@sacids.org

About Catholic University of Health and Allied Sciences-Bugando

The Catholic university of Health and Allied Sciences is accredited higher learning institution owned by Tanzania Episcopal Conference. It was established under University Act in 2003. Its mission is to provide skilled and competent human resources in the health sector that is vested with moral and ethical values; Search, discover and communicate the truth to advance the frontiers of knowledge and provide quality services to the community.
Website: <http://www.bugando.ac.tz>